



For Professional Use Only


Sacace Molecular Genetics

Handbook

Real Time PCR kits for detection of Single Nucleotide Polymorphisms (SNPs)

REF see below kits table

 **60**

 **96**

NAME

Sacace Molecular Genetics

INTRODUCTION

A single nucleotide polymorphism (SNP pronounced "snip") is a DNA polymorphisms at the level of a single nucleotide, a single base mutation in DNA. SNPs are 'conserved' across the genome and represent the most simple form and most common source of genetic polymorphism in the human genome: 90% of all human DNA polymorphisms are associated with SNPs and variation frequency is about 1 every 1000bp in the human genome (Sachidanandam et al.,2001).

The SNPs in the genome can affect gene functions, protein structure or expression and for these reasons they are used as markers in genetic disease studies (Kim & Mishra, 2007).

It's sometimes possible to correlate a SNP with a particular trait or disease: susceptibility to disease may also be described as an 'unfortunate trait' that can be assessed checking if the mutated (unfortunate) polymorphism is carried in both alleles.

SNPs testing can be applied to:

- Diagnostics / risk profiling
- Drug response prediction
- Gene function identification

Several SNPs have been associated to genetic susceptibility to different diseases and disorders like for example:

- Hypertension
- Fibrinolysis
- Myocardial infarction
- Ischemic stroke
- Cancer
- Metabolic disorders

In order to perform SNP genotyping, two specific probes labeled with different dyes are used, the first for the wild type allele and the second for the mutant allele. If the assay results in the emission of only the first fluorescent color, then the individual is homozygous wild type at that locus. If the assay results in the emission of only the second fluorescent color, then the individual is homozygous mutant. And finally, if both fluorescent colors are produced, then the individual is heterozygous.

INTENDED USE

Sacace Molecular Genetics Kits are intended for detection and allelic discrimination of genetic polymorphisms associated with inherited susceptibility to increased risk of disease, or to different response to drugs.

PRINCIPLE OF ASSAY

Sacace Molecular Genetics Kits are qualitative tests that allow the detection by Real Time PCR based on the amplification of the genome specific region using specific primers. In Real Time PCR the amplified product is detected using fluorescent dyes. These dyes are linked to oligonucleotide probes that bind specifically to the amplified product. The real-time monitoring of the fluorescence intensities during the reaction allows the detection of accumulating product without re-opening of the reaction tubes after the PCR run.

MATERIALS PROVIDED

Option No.1: Ready to use 0,2 ml tube format (TXXXXX-50-T)

- **60 ready to use 0,2 ml PCR tubes** (each PCR tube contains 15 µl of PCR mix)
- **Taq polymerase**, 0,3 ml (1 vial)
- **Negative control C-**, 0,1 mL (1 vial)
- **C+ Homozygous Wild Type (allele 1-1)**, 50 µL (1 vial)
- **C+ Heterozygous (allele 1-2)**, 50 µL (1 vial)
- **C+ Homozygous Mutant (allele 2-2)**, 50 µL (1 vial)

Contains reagents for 60 tests.

Option No.2: Ready to use 12x8 strip format (TXXXXX-96-S)

- **12 x 8 strip ready to use** (each PCR tube contains 15 µl of PCR mix)
- **Taq polymerase**, 0,5 ml (1 vial)
- **Negative control C-**, 0,1 mL (1 vial)
- **C+ Homozygous Wild Type (allele 1-1)**, 50 µL (1 vial)
- **C+ Heterozygous (allele 1-2)**, 50 µL (1 vial)
- **C+ Homozygous Mutant (allele 2-2)**, 50 µL (1 vial)

Contains reagents for 96 tests.

KITS TABLE

Code	Gene	Polymorphism details	Fluorescence Channel / Allele
T01273	MTHFR	Glu429Ala 1298 A>C Rs1801131	HEX: Glu (A) – allele 1
			FAM: Ala (C) – allele 2
T01105	F7	Arg 353 Gln C <u>G</u> G 353 C <u>A</u> G rs6046	HEX: Arg (G) – allele 1
			FAM: Gln (A) – allele 2
T01106	ITGB3	Leu 33 Pro C <u>T</u> G 33 C <u>C</u> G rs5918	HEX: Leu (T) – allele 1
			FAM: Pro (C) – allele 2
T01107	FGB	G-455A rs1800790	HEX: G – allele 1
			FAM: A – allele 2
T01120	PAI	-675 5G/4G rs1799768	HEX: 5G – allele 1
			FAM: 4G – allele 2
T01124	MTRR	Ile 22 Met ATA 22 ATG A 66 G rs1801394	HEX: Ile (A) – allele 1
			FAM: Met (G) – allele 2
T01143	MTR	Asp 919 Gly G <u>A</u> C 919 G <u>G</u> C rs1805087	HEX: Asp (A) – allele 1
			FAM: Gly (G) – allele 2
T01155	ITGA2	Phe 224 Phe T <u>T</u> C 224 T <u>T</u> I C807T rs1126643	HEX: C – allele 1
			FAM: T – allele 2
T01179	GPIBA	Thr 145 Met C482T A <u>C</u> G 145 A <u>I</u> G rs6065	HEX: Thr (C) – allele 1
			FAM: Met (T) – allele 2
T01354	GPIBA	T -5 C rs2243093	HEX: T – allele 1
			FAM: C – allele 2
T01355	F13A1	Val35Leu G <u>T</u> G 35 <u>I</u> TG rs5985	HEX: Val (G) – allele 1
			FAM: Leu (T) – allele 2
T01356	F12	C -4 T rs1801020	HEX: C – allele 1
			FAM: T – allele 2
T01329	FTO *	A 23525 T rs9939609	HEX: A – allele 1
			FAM: T - allele 2
T01331	CYP3A5 *	G 6986 A rs776746	HEX: G – allele 1
			FAM: A – allele 2
T01182	NOS3 *	C786T	HEX: C – allele 1
			FAM: T – allele 2

Code	Gene	Polymorphism details	Fluorescence Channel / allele
T01357	SELPLG	Met 62 Ile AT <u>G</u> 62 AT <u>A</u> rs2228315	HEX: Met (G) – allele 1
			FAM: Ile (A) – allele 2
T01104	CYP2C9	Arg 144 Cys <u>C</u> GT 144 <u>I</u> GT rs1799853	HEX: Arg (C) – allele 1
			FAM: Cys (T) – allele 2
T01111	CYP2C9	Ile 359 Leu <u>A</u> TT 359 <u>C</u> TT rs1057910	HEX: Ile (A) – allele 1
			FAM: Leu (C) – allele 2
T01144	VKORC1	C1173T rs9934438	HEX: C – allele 1
			FAM: T – allele 2
T01145	VKORC1	G3730A rs7294	HEX: G – allele 1
			FAM: A – allele 2
T01335	PPARG2	Pro12Ala <u>C</u> CA 12 <u>G</u> CA rs1801282	HEX: C – allele 1
			FAM: G – allele 2
T01358	ADRB2	Gln27Glu rs1042714	HEX: C – allele 1
			FAM: G – allele 2
T01359	ADRB2	Arg16Gly rs1042713	HEX: T – allele 1
			FAM: C – allele 2
T01360	ADRB3	Trp64Arg rs4994	HEX: A – allele 1
			FAM: G – allele 2
T01361	FABP2	Ala54Thr rs1799883	HEX: G – allele 1
			FAM: A – allele 2
T01349	IL28B	rs8099917 T>G	HEX: T – allele 1
			FAM: G – allele 2
T01371	IL28B	rs12979860 C>T	HEX: C – allele 1
			FAM: T – allele 2
T01303	SLCO1B1	Val 174 Ala GTG 521 GCG rs 4149056	HEX: Val(T) – allele 1
			FAM: Ala (C) – allele 2

Code	Gene	Polymorphism details	Fluorescence Channel / allele
T01118	AGT (1)	Thr174Met rs4762	HEX: Thr(C) – allele 1
			FAM: Met(T) – allele 2
T01119	AGT (2)	Met235Thr Rs699	HEX: Met(T) – allele 1
			FAM: Thr(C) – allele 2
T01131	AGTR1	A1166C Rs51186	HEX: A – allele 1
			FAM: C – allele 2
T01148	ApoE	Leu28Pro rs769452	HEX: Leu(T) – allele 1
			FAM: Pro(C) – allele 2
T01149	LPL	Ser474Ter Rs328	HEX: Ser(C) – allele 1
			FAM: Ter(G) – allele 2
T01323	CYP2C19(*2)	G681A rs4244285	HEX: G – allele 1
			FAM: A – allele 2
T01324	CYP2C19(*3)	Trp212Ter TGG 212 TGA rs4986893	HEX: Trp(G) – allele 1
			FAM: Ter(A) – allele 2
T01171	IL17A	IL17A G-197A	HEX: G – allele 1
			FAM: A – allele 2
T01352	COMT	Val158Met rs4680	HEX: Val (G) – allele 1
			FAM: Met (A) – allele 2
T01272	ACE	Alu Ins/Del rs4646994	HEX: Ins – allele 1
			FAM: Del – allele 2
T01177	TNF	G-308A rs1800629	HEX: G – allele 1
			FAM: A – allele 2

** in those kits Allele 1 on HEX identifies the “mutant” genotype and Allele 2 on FAM identifies the “wild type” genotype: always refer to the allele base name in the results (for example CC or CG or GG); refer to the reference rs number (for example rs12979860) as indicated in the “kits table” for details about the polymorphism.*

MATERIALS REQUIRED BUT NOT PROVIDED

Zone 1: sample preparation

- DNA extraction kit
- Biological cabinet
- Desktop microcentrifuge for “eppendorf” type tubes
- Dry heat block
- Vortex mixer
- Pipettes
- Sterile pipette tips with filters
- 1,5 ml polypropylene sterile tubes
- Biohazard waste container
- Refrigerator, Freezer

Zone 2: Real Time amplification

- Real Time Thermal cycler
- Workstation
- Pipettes (adjustable)
- Sterile pipette tips with filters
- Vortex mixer
- Desktop centrifuge with rotor for 1,5/2,0 ml tubes
- Freezer, refrigerator
- Tube racks

STORAGE INSTRUCTIONS

Sacace Molecular Genetics kits must be stored at 2-8°C. The kits can be shipped at 2-8°C and stored as indicated immediately on receipt.

STABILITY

Sacace Molecular Genetics kits are stable up to the expiration date indicated on the kit label. The product will maintain performance through the control date printed on the label. Exposure to light, heat or humidity may affect the shelf life of some of the kit components and should be avoided. Repeated thawing and freezing of these reagents should be avoided, as this may reduce the sensitivity. Components stored under conditions other than those stated on the labels may not perform properly and may adversely affect the assay results.

QUALITY CONTROL

In accordance with Sacace's ISO 13485-Certified Quality Management System, each lot is tested against predetermined specifications to ensure consistent product quality.

WARNINGS AND PRECAUTIONS



For Research Use Only

The user should always pay attention to the following:

- Use sterile pipette tips with aerosol barriers and use new tip for every procedure.
- Store extracted positive material (samples, controls and amplicons) away from all other reagents and add it to the reaction mix in a separate area.
- Use disposable gloves, laboratory coats and eye protection when handling specimens and reagents. Thoroughly wash hands afterwards.
- Do not eat, drink, smoke, apply cosmetics, or handle contact lenses in laboratory work areas.
- Do not use a kit after its expiration date.
- Dispose of all specimens and unused reagents in accordance with local authorities' regulations.
- Specimens should be considered potentially infectious and handled in a biological cabinet in accordance with appropriate biosafety practices.
- Clean and disinfect all sample or reagent spills using a disinfectant such as 0.5% sodium hypochlorite, or other suitable disinfectant.
- Avoid sample or reagent contact with the skin, eyes, and mucous membranes. If skin, eyes, or mucous membranes come into contact, rinse immediately with water and seek medical advice immediately.
- Material Safety Data Sheets (MSDS) are available on request.
- Use of this product should be limited to personnel trained in the techniques of DNA amplification.
- The laboratory process must be one-directional, it should begin in the Extraction Area and then move to the Amplification and Detection Areas. Do not return samples, equipment and reagents to the area in which the previous step was performed.

PRODUCT USE LIMITATIONS

Use of this product should be limited to personnel trained in the techniques of DNA amplification (UNI EN ISO 18113-2:2012). Strict compliance with the user manual is required for optimal PCR results. Attention should be paid to expiration dates printed on the box and labels of all components. Do not use a kit after its expiration date.

SAMPLE COLLECTION, STORAGE AND TRANSPORT

Sacace Molecular Genetics Kits can analyze genomic DNA extracted from:

- *whole blood* collected in EDTA tubes;
- *Buccal swab*: insert the swab into the nuclease-free 1,5 ml tube and add 0,2 ml of Transport medium. Vigorously agitate swabs in medium for 15-20 sec.

Specimens can be stored at +2-8°C for no longer than 24 hours, or freeze at -20°C to -80°C.

Transportation of clinical specimens must comply with country, federal, state and local regulations for the transport of etiologic agents.

DNA ISOLATION

The following isolation kits are recommended:

- ⇒ **Genomic column DNA Express** – spin column extraction kit (Sacace, [REF](#) K-1-1/E)
- ⇒ **SaMag Blood DNA extraction kit** (Sacace, [REF](#) SM001);
- ⇒ **QIAamp DNA Blood mini kit** (Qiagen, [REF](#) 51104);
- ⇒ **DNA-Sorb-A** (Sacace, REF K-1-1/A) for buccal swab;

Please carry out DNA extraction according to the manufacturer's instruction.

PROTOCOL

Sacace Molecular Genetics kits do not include reagents required for sample preparation and DNA extraction. Blood samples and biological materials must be processed by using the recommended kits or those with similar performances of quality and quantity of extracted DNA. Use of blood samples collected in tubes containing heparin is not recommended.

The analysis of the genomic DNA specimens using **Sacace Molecular Genetics** kits includes the following stages:

1. Preparing the Real Time PCR;
2. Real Time PCR analysis;
3. Data analysis with the software of Real Time PCR instrument;
4. Results analysis and conclusions.

EXPERIMENTAL PROTOCOL

Total reaction volume: 25 µl

1. Prepare the necessary number of ready-to-use PCR tubes (samples + 3 pos controls + 1 neg control).
2. Spin for 3-5 sec the **Taq polymerase**, mix by pipetting and **add 5 µl** to each PCR tube.
3. Add into the corresponding PCR tubes **5.0 µl** of extracted DNA from sample:
 - **DNA sample**Add into the corresponding PCR tubes **5.0 µl** of controls:
 - **C+ Homozygous (allele 1-1)**
 - **C+ Heterozygous (allele 1-2)**
 - **C+ Homozygous (allele 2-2)**
 - **Negative Control C-**
4. Spin the tubes for 3–5 seconds to collect the drops.
5. Insert the tubes in the Real-time PCR instrument.

Amplification

Create a temperature profile on your instrument as follows:

Step	Plate or modular type instruments ¹			Rotor type instruments ²		
	Temperature, °C	Time	Cycles	Temperature, °C	Time	Cycles
Hold	80	2 min	1	80	2 min	1
Hold	94	3 min	1	95	3 min	1
Cycling	94	15 s	5	95	10 s	40
	64	40 s		60	40 s fluorescence detection	
Cycling 2	94	15 s	35			
	64	40 s fluorescence detection				

¹ For example, SaCycler-96™ (Sacace); CFX-96 / iQ5™ (BioRad); Mx3005P™ (Agilent); ABI® 7500 Real Time PCR (Applied)*; LightCycler® 96 (Roche).

² For example Rotor-Gene™ 6000/Q (Corbett Research, Qiagen); when using Rotor Gene instrument the tube caps may be marked and it's recommended to cut the strip into two equal parts (4 tubes each)

* To perform the test with ABI 7500 (Applied) a disposable adapter provided with the kit has to be used. Additional adapters can be purchased separately.

Fluorescence is detected in FAM/Green, JOE/Yellow/HEX fluorescence channels.

DATA ANALYSIS

The fluorescent signal intensity is detected in 2 channels as shown in the table below:

FAM	HEX
Allele 2 ("mutant"*)	Allele 1 ("wild type"*)

Note: Please refer to the "Kits Table" at the beginning of this manual to check the details of the nucleotides substitution for each polymorphism.

* in some kits Allele 1 on HEX identifies the "mutant" genotype and Allele 2 on FAM identifies the "wild type" genotype: always refer to the "kits table" and the rs code at the beginning of the manual for details about the polymorphism.

Interpretation of results for Rotorgene 6000/Q (Corbett Research, Qiagen):

Principle of interpretation:

- Signal only in allele 1 (Yellow) : homozygous "wild type"
- Signal only in allele 2 (Green) : homozygous "mutant"
- Signal in both allele 1 and allele 2 : heterozygous

Genotyping		
Genotype	Reacting Channels	Reacting Channels
Allele 2	Cycling A.Green	
Heterozygous	Cycling A.Green	Cycling A.Yellow
Allele 1		Cycling A.Yellow

Click **Analysis**, click **Other**, select **Allelic Discrimination**, select **Slope Correct**, click **Eliminate cycles before / Ignore first** and insert value. Insert the **Threshold** and **Outlier removal** values as in the following table:

Code	Gene	Polymorphism	Channel / allele	Threshold	Slope Correct	Outlier Removal	Ignore first
T01105	F7	Arg 353 Gln CGG 353 CAG rs6046	Yellow: Arg (G)	0,03	On	10%	10
			Green: Gln (A)				
T01358	ADRB2	Gln27Glu rs1042714	Yellow: C	0,03	On	15%	5
			Green: G				
T01359	ADRB2	Arg16Gly rs1042713	Yellow: T	0,03	On	15%	5
			Green: C				
T01360	ADRB3	Trp64Arg rs4994	Yellow: A	0,03	On	15%	5
			Green: G				
T01349	IL28B	rs8099917 T>G	Yellow: T	0,03	On	15%	5
			Green: G				
T01371	IL28B	rs12979860 C>T	Yellow: C	0,03	On	0%	0
			Green: T				
T01106	ITGB3	Leu 33 Pro CTG 33 CCG rs5918	Yellow: Leu (T)	0,1	On	5%	5
			Green: Pro (C)				
T01107	FGB	G-455A rs1800790	Yellow: G	0,03	On	10%	10
			Green: A				
T01120	PAI	-675 5G/4G rs1799768	Yellow: 5G	0,15	On	10%	5
			Green: 4G				

Code	Gene	Polymorphism	Channel/allele	Threshold	Slope Correct	Outlier Removal	Ignore first
T01143	MTR	Asp 919 Gly GAC 919 GGC rs1805087	Yellow: Asp(A)	0,03	On	15%	5
			Green: Gly (G)				
T01124	MTRR	Ile 22 Met ATA 22 ATG A 66 G rs1801394	Yellow: Ile (A)	0,03	On	15%	5
			Green: Met(G)				
T01155	ITGA2	Phe 224 Phe TTC 224 TTT C807T rs1126643	Yellow: C	0,03	On	10%	10
			Green: T				
T01179	GPIBA	Thr 145 Met ACG 145 ATG rs6065	Yellow: Thr (C)	0,03	On	15%	5
			Green: Met (T)				
T01354	GPIBA	T -5 C rs2243093	Yellow: Thr (C)	0,03	On	15%	5
			Green: Met (T)				
T01273	MTHFR	Glu429Ala 1298 A>C	Yellow: Glu (A)	0,03	On	15%	5
			Green: Ala (C)				
T01355	F13A1	Val35Leu GTG 35 TTG rs5985	Yellow: Val (G)	0,03	On	10%	10
			Green: Leu (T)				
T01356	F12	C -4 T rs1801020	Yellow: C	0,03	On	15%	5
			Green: T				
T01329	FTO	A 23525 T rs9939609	Yellow: A	0,03	On	10%	5
			Green: T				
T01331	CYP3A5	G 6986 A rs776746	Yellow: G	0,015	On	0%	5
			Green: A				
T01104	CYP2C9	Arg 144 Cys CGT 144 TGT rs1799853	Yellow: Arg(C)	0,03	On	20%	5
			Green: Cys(T)				
T01111	CYP2C9	Ile 359 Leu ATT 359 CTT rs1057910	Yellow: Ile(A)	0,03	On	20%	15
			Green: Leu(C)				
T01144	VKORC1	C1173T rs9934438	Yellow: C	0,03	On	0%	0
			Green: T				
T01145	VKORC1	G3730A rs7294	Yellow: G	0,03	On	10%	0
			Green: A				
T01335	PPARG2	Pro12Ala CCA 12 GCA rs1801282	Yellow: C	0,03	On	15%	5
			Green: G				
T01361	FABP2	Ala54Thr rs1799883	Yellow: G	0,03	On	15%	5
			Green: A				
T01303	SLCO1B1	Val 174 Ala GTG 521 GCG rs 4149056	Yellow: Val(T)	0,03	On	10%	5
			Green: Ala(C)				
T01182	NOS3	C786T	Yellow: C	0,05	On	10%	5
			Green: T				
T01118	AGT (1)	Thr174Met rs4762	Yellow: Thr(C)	0,03	On	10%	5
			Green: Met(T)				
T01119	AGT (2)	Met235Thr Rs699	Yellow: Met(T)	0,03	On	10%	5
			Green: Thr(C)				
T01131	AGTR1	A1166C Rs51186	Yellow: A	0,03	On	10%	5
			Green: C				
T01148	ApoE	Leu28Pro	Yellow: Leu(T)	0,03	On	10%	5
			Green: Pro(C)				
T01149	LPL	Ser447Ter	Yellow: Ser(C)	0,03	On	10%	5
			Green: Ter(G)				
T01323	CYP2C19(*2)	G681A rs4244285	Yellow: G	0,06	On	15%	10
			Green: A				
T01324	CYP2C19(*3)	Trp212Ter TGG 212 TGA rs4986893	Yellow: Trp (G)	0,06	On	15%	10
			Green: Ter (A)				
T01171	IL17A	IL17A G-197A	Yellow: G	0,06	On	15%	0
			Green: A				

T01352	COMT	Val158Met rs4680	Yellow: Val(G)	0,03	On	15%	0
			Green: Met (A)				
T01272	ACE	Alu Ins/Del rs4646994	Yellow: Ins	0,03	On	10%	5
			Green: Del				
t01177	TNF	G-308A rs1800629	Yellow: G	0,03	on	15%	0
			Green: A				

NOTE for Rotorgene 6000/Q (Corbett Research, Qiagen): if a Ct value is higher than 37 the sample is considered invalid and must be repeated. If there is no Ct value in both channels the sample is invalid and must be repeated starting from the extraction.

Interpretation of results for CFX-96/iQ5 (Bio-rad):

Principle of interpretation:

- **Signal only in allele 1 (channel HEX) : homozygous “wild type”**
- **Signal only in allele 2 (channel FAM) : homozygous “mutant”**
- **Signal in both allele 1 and allele 2 (channels HEX and FAM) : heterozygous**

Set **Baseline Cycles** at 5-15 and **Crossing Threshold** value at 100.

NOTE FOR CFX-96/iQ5 (Bio-rad): if a Ct value is higher than 32 the sample is considered invalid and must be repeated. If there is no Ct value in both channels the sample is invalid and must be repeated starting from the extraction.

Interpretation of results for SaCycler-96 (Sacace Biotechnologies):

Principle of interpretation:

- **Signal only in allele 1 (channel HEX) :** homozygous “*wild type*”
- **Signal only in allele 2 (channel FAM) :** homozygous “*mutant*”
- **Signal in both allele 1 and allele 2 (channels HEX and FAM) :** heterozygous

NOTE: when creating new test for Sacace Molecular Genetics, select “**Analysis of polymorphisms (two probes)**”, name “a” on FAM channel and name “b” on HEX channel. Set **Heterozygote dCp < 3,0** and **Homozygote dCp > 6** (see pictures below).

1. Analysis
 Type: Analysis of polymorphisms (two probes) ▼
 Method: dF/dT ▼

5. Mixture volume: 35 mcL

6. Fluorofors:
 Fam ▼ Hex ▼ Rox ▼ Cy5 ▼ Cy5.5 ▼
 a ▼ b ▼ is absent ▼ is absent ▼ is absent ▼


7. Polimorphisms analysis criterion:
 Heteozygote dCp < 3,0
 Homozygote dCp > 6,0


To analyse results, be sure to select “**Analysis of polymorphisms (two probes)**” as Analysis type and “**Curve Shape (Cp)**” as Method.

	N°	Identifier	R	Test	Tube type	Concentration				
						Fam	Hex	Rox	Cy5	Cy5.5
A1	1	Sample_1_f2	<input type="checkbox"/>	snp_new	<input type="checkbox"/>	b	a	-	-	-
A2	2	Sample_2	<input type="checkbox"/>	snp_new	<input type="checkbox"/>	b	a	-	-	-
A3	3	Sample_3	<input type="checkbox"/>	snp_new	<input type="checkbox"/>	b	a	-	-	-
A4	4	Sample_4	<input type="checkbox"/>	snp_new	<input type="checkbox"/>	b	a	-	-	-
A5	5	pos 1-1 (wt)	<input type="checkbox"/>	snp_new	C+	b	a	-	-	-
A6	6	pos 1-2 (het)	<input type="checkbox"/>	snp_new	C+	b	a	-	-	-
A7	7	pos 2-2 (mut)	<input type="checkbox"/>	snp_new	C+	b	a	-	-	-
A8	8	C_-	<input type="checkbox"/>	snp_new	C-	b	a	-	-	-

Analysis type: Analysis of polymorphisms (two probes) ▼

Method: Curve Shape (Cp) ▼

Click on the icon for changing the parameter of data analysis  , a new window will show up.



Parameters of the analysis

1. Criterion of the positive PCR result: 90 %

2. Threshold value 10 StD at background part of PCR curve

3. Criteria to validity result

bottom edge/threshold of the positive result 10 % F(Cp)

top edge/threshold to normalizations data 10 % F(Cp)

4. Normalization data

5. Parameters of polymorphisms analysis:

heterozygote dCp < 3,0

homozygote dCp > 6,0

Additional Apply Cancel

Set **90%** as “*Criterion of the positive PCR result*”; “*Normalization data*” checkbox must be **deselected**.

Select checkbox “**Criteria to validity result**” and insert between **10-20% F(Cp)** for “*bottom edge/threshold of the positive result*” and insert between **10-20% F(Cp)** for “*top edge/threshold to normalizations data*”, then click “**Apply**”.

The results will be displayed in the table on the right (see below pictures as reference).

Example of results:

Results		Statistics				
N	Identificator	Polimorphism		dCp	Cp Fam	Cp Hex
A1	Sample_1_f2	a	b	0,2	19,2	19,1
A2	Sample_2	b	b	>17		17,7
A3	Sample_3	b	b	>16		18,3
A4	Sample_4	b	b	>16		18,8
A5	K+	b	b	>15		19,4
A6	K+	a	b	0,1	18,9	19,0
A7	K+	a	a	>16		18,4
A8	C_-	-	-			

D1	Sample_1_mthfr	a	b	0,1	18,6	18,7
D2	Sample_2	a	a	>18	17,0	
D3	Sample_3	b	b	>17		17,5
D4	Sample_4	b	b	>16		18,1
D5	K+	b	b	>19		15,7
D6	K+	a	b	0,4	13,9	13,5
D7	K+	a	a	>21		13,0
D8	C_-	-	-			

a = FAM (allele2) b = HEX (allele1)

a b = sample heterozygous (both allele 1 and allele 2 present)

b b = sample homozygous with only allele 1 present (“wild type”)

a a = sample homozygous with only allele 2 present (“mutant”)

NOTE FOR SaCycler-96 (Sacace Biotechnologies): if a Ct value is higher than 32 the sample is considered invalid and must be repeated. If there is no Ct value in both channels the sample is invalid and must be repeated starting from the extraction.

KEY TO SYMBOLS USED



List Number



For Research Use Only



Expiration Date



Store at



Manufacturer



Consult instructions for use



Lot Number



Caution!



Contains sufficient
for <n> tests



Version

NCA

Negative Control of
Amplification

NCE

Negative control of
Extraction

IC

Internal Control

* SaCycler™ is a registered trademark of Sacace Biotechnologies

* iQ5™ is a registered trademark of Bio-Rad Laboratories

* Rotor-Gene™ Technology is a registered trademark of Qiagen

* MX3005P® is a registered trademark of Agilent Technologies

* ABI® is a registered trademark of Applied Biosystems

* LightCycler® 96 is trademark of Roche



Sacace Biotechnologies Srl

via Scalabrini, 44 – 22100 – Como – Italy Tel +390314892927 Fax +390314892926

mail: info@sacace.com web: www.sacace.com